

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 February 2002 (07.02.2002)

PCT

(10) International Publication Number
WO 02/10166 A1

(51) International Patent Classification⁷: **C07D 471/16**,
A61K 31/495, C07D 471/14

(21) International Application Number: PCT/US01/21678

(22) International Filing Date: 9 July 2001 (09.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/222,451 2 August 2000 (02.08.2000) US

(71) Applicant (for all designated States except US): **LILLY ICOS LLC** [US/US]; 1209 Orange Street, Wilmington, DE 19801 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ORME, Mark, W.** [US/US]; 4235 Francis Avenue #203, Seattle, WA 98103 (US). **SAWYER, Jason, Scott** [US/US]; 5718 North Winthrop Avenue, Indianapolis, IN 46220 (US). **SCHULTZE, Lisa, M.** [US/US]; 16110 N.E. 175th Street, Woodinville, WA 98072 (US).

(74) Agent: **NAPOLI, James, J.**; Marshall, O'Toole, Gerstein, Murray & Borun, 6300 Sears Tower, 233 South Wacker Drive, Chicago, IL 60606 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: FUSED HETEROCYCLIC DERIVATIVES AS PHOSPHODIESTERASE INHIBITORS




(57) Abstract: Compounds of general structural formula (I) and use of the compounds and salts and solvates thereof, as therapeutic agents.

FUSED HETEROCYCLIC DERIVATIVES AS PHOSPHODIESTERASE INHIBITORS

5 This invention relates to a series of compounds, to methods of preparing the compounds, to pharmaceutical compositions containing the compounds, and to their use as therapeutic agents. In particular, the invention relates to compounds that
10 are potent and selective inhibitors of cyclic guanosine 3',5'-monophosphate specific phosphodiesterase (cGMP-specific PDE), in particular PDE5, and have utility in a variety of therapeutic areas wherein such inhibition is considered beneficial, including
15 the treatment of cardiovascular disorders and erectile dysfunction.

SUMMARY OF THE INVENTION

20 The present invention provides compounds of formula (I)




(I)

30 wherein R⁰, independently, is selected from the group consisting of halo and C₁₋₆alkyl;

- 2 -

R^1 is selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, $haloC_{1-6}$ alkyl, C_{3-8} cycloalkyl, C_{3-8} cycloalkyl C_{1-3} alkyl, aryl C_{1-3} alkyl, and heteroaryl C_{1-3} alkyl;

5 R^2 is selected from the group consisting of an optionally substituted monocyclic aromatic ring selected from the group consisting of benzene, thiophene, furan, and pyridine, and an optionally substituted bicyclic ring



10 wherein the fused ring A is a 5- or 6-membered ring, saturated or partially or fully unsaturated, and comprises carbon atoms and optionally one or two heteroatoms selected from oxygen, sulfur, and
15 nitrogen;

20 X is selected from the group consisting of NR^a , O, S, CR^3R^4 , $CH_2CR^3R^5$, and $CR^3R^5CH_2$;

R^3 is selected from the group consisting of hydrogen and C_{1-3} alkyl;

25 or R^1 and R^3 together represent a 3- or 4-membered alkyl or alkenyl chain component of a 5- or 6-membered ring,

or NR^a , R^1 , and the nitrogen to which R^1 is attached, form a 5- or 6-membered heterocyclic ring;

30 R^4 is selected from the group consisting of C_{1-6} alkyl, $haloC_{1-4}$ alkyl, aryl, heteroaryl, Het, C_{3-8} cycloalkyl, OR^a , $C(=O)OR^a$, $C(=O)R^a$, $C(=O)NR^aSO_2R^b$, $C(=O)NR^aR^b$, $C(=S)NR^aR^b$, C_{1-6} alkylene $C(=O)OR^a$, C_{1-4} alkyl-

- 3 -

eneC(=O)NR^aR^b, C₁₋₄alkyleneNR^aR^b, C₁₋₄alkyleneOR^a, C₁₋₄-
alkyleneSO₂NR^aR^b, C₁₋₄alkylenearyl, C₁₋₄alkylenehetero-
oaryl, C₁₋₄alkyleneHet, C₁₋₄alkyleneC(=O)C₁₋₄alkylene-
Het, C₁₋₄alkyleneC(=O)Het, C₁₋₄alkyleneC(=O)OR^a, C(=O)-
5 C₁₋₄alkyleneHet, C(=O)NR^aalkyleneOR^b, C(=O)NR^aC₁₋₄alkyl-
eneHet, NR^aC₁₋₄alkyleneNR^aR^b, NR^aC(=O)R^b, NR^aC(=O)NR^aR^b,
N(SO₂C₁₋₄alkyl)₂, and NR^a(SO₂C₁₋₄alkyl), with the pro-
viso that when R³ is hydrogen, R⁴ is different from
C₁₋₃alkyl;

10 or R¹ and R⁴ together represent a 3- or 4-
membered carbocyclic or heterocyclic chain compo-
nent, either saturated or unsaturated, of a 5- or 6-
membered ring;

15 or R³ and R⁴ together represent a 4- to 6-
membered alkyl or alkenyl chain component of a 5- to
7-membered ring;

R⁵ is selected from the group consisting of
hydrogen, C₁₋₆alkyl, haloC₁₋₄alkyl, aryl, heteroaryl,
Het, C₃₋₈cycloalkyl, OR^a, C(=O)OR^a, C(=O)R^a, C(=O)-
20 NR^aSO₂R^b, C(=O)NR^aR^b, C(=S)NR^aR^b, C₁₋₆alkyleneC(=O)OR^a,
C₁₋₄alkyleneC(=O)NR^aR^b, C₁₋₄alkyleneNR^aR^b, C₁₋₄alkylene-
OR^a, C₁₋₄alkyleneSO₂NR^aR^b, C₁₋₄alkylenearyl, C₁₋₄alkyl-
eneheteroaryl, C₁₋₄alkyleneHet, C₁₋₄alkyleneC(=O)C₁₋₄-
alkyleneHet, C₁₋₄alkyleneC(=O)Het, C₁₋₄alkyleneC(=O)-
25 OR^a, C(=O)C₁₋₄alkyleneHet, C(=O)NR^aalkyleneOR^b, C(=O)-
NR^aC₁₋₄alkyleneHet, NR^aC₁₋₄alkyleneNR^aR^b, NR^aC(=O)R^b,
NR^aC(=O)NR^aR^b, N(SO₂C₁₋₄alkyl)₂, and NR^a(SO₂C₁₋₄alkyl);

R^a and R^b, independently, are selected from
the group consisting of hydrogen, C₁₋₈alkyl, aryl,
30 heteroaryl, haloC₁₋₆alkyl, C₁₋₄alkyleneHet, arylC₁₋₃-
alkyl, heteroarylC₁₋₃alkyl, C₁₋₄alkylenearyl, C₃₋₈-
cycloalkyl, C₁₋₃alkyleneheteroaryl, and Het;

- 4 -

Het is a 4- to 7-membered heterocyclic group, saturated or partially unsaturated, containing at least one heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur, and optionally substituted with one or more of C₁₋₄alkyl, NR^aR^b, and C(=O)OR^a;

q is 0, 1, 2, 3, or 4; and
pharmaceutically acceptable salts and solvates (e.g., hydrates) thereof.

As used herein, the term "alkyl" includes straight chained and branched hydrocarbon groups containing the indicated number of carbon atoms, typically methyl, ethyl, and straight chain and branched propyl and butyl groups. The hydrocarbon group can contain up to 16 carbon atoms. The term "alkyl" includes "bridged alkyl," i.e., a C₆-C₁₆ bicyclic or polycyclic hydrocarbon group, for example, norbornyl, adamantyl, bicyclo[2.2.2]octyl, bicyclo[2.2.1]heptyl, bicyclo[3.2.1]octyl, or decahydronaphthyl. The term "cycloalkyl" is defined as a cyclic C₃-C₈ hydrocarbon group, cyclopropyl, cyclobutyl, cyclohexyl, and cyclopentyl. The terms "alkenyl" and "alkynyl" are defined similarly to the term "alkyl," except the hydrocarbon group contains a carbon-carbon double bond or carbon-carbon triple bond, respectively.

The term "alkylene" refers to an alkyl group having a substituent. For example, the term "C₁₋₃alkylenearyl" refers to an alkyl group containing one to three carbon atoms and substituted with an aryl group. The term "alkenylene" as used herein is similarly defined, and contains the indicated number of carbon atoms and a carbon-carbon double

- 5 -

bond, and includes straight chained and branched alkenylene groups, like ethylenylene.

5 The term "halo" or "halogen" is defined herein to include fluorine, bromine, chlorine, and iodine.

10 The term "haloalkyl" is defined herein as an alkyl group substituted with one or more halo substituents, either fluoro, chloro, bromo, iodo, or combinations thereof. Similarly, "halocycloalkyl" is defined as a cycloalkyl group having one or more halo substituents.

15 The term "aryl," alone or in combination, is defined herein as a monocyclic or polycyclic aromatic group, preferably a monocyclic or bicyclic aromatic group, e.g., phenyl or naphthyl, that can be unsubstituted or substituted, for example, with one or more, and in particular one to three, hydroxy, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkyl, nitro, amino, carboxy (CO_2H), CO_2R^a , alkylamino, acylamino, alkylthio, alkylsulfinyl, and alkylsulfonfyl. Exemplary aryl groups include phenyl, naphthyl, tetrahydronaphthyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-methylphenyl, 4-methoxyphenyl, 3-trifluoromethylphenyl, 4-nitrophenyl, and the like. The terms "aryl C_{1-3} alkyl" and "heteroaryl C_{1-3} alkyl" are defined as an aryl or heteroaryl group having a C_{1-3} alkyl substituent.

30 The term "heteroaryl" is defined herein as a monocyclic or bicyclic ring system containing one or two aromatic rings and containing at least one nitrogen, oxygen, or sulfur atom in an aromatic ring, and which can be unsubstituted or substituted, for example, with one or more, and in particular one

- 6 -

to three, substituents, like halo, alkyl, hydroxy, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkyl, nitro, amino, carboxy (CO_2H), CO_2R^a , alkylamino, acylamino, alkylthio, alkylsulfinyl, and alkylsulfonyl. Non-limiting examples of heteroaryl groups include thienyl, furyl, pyridyl, oxazolyl, quinolyl, isoquinolyl, indolyl, triazolyl, tetrazolyl, isothiazolyl, isoxazolyl, imidazolyl, benzothiazolyl, pyrazinyl, pyrimidinyl, thiazolyl, and thiadiazolyl.

The term "Het" includes a 4- to 7-membered heterocycloalkyl group, including, but not limited to, morpholinyl, piperidyl, pyrrolidinyl, or piperazinyl.

The term "alkoxy" is defined as -OR, wherein R is alkyl.

The term "alkoxyalkyl" is defined as an alkyl group wherein a hydrogen has been replaced by an alkoxy group. The term "(alkylthio)alkyl" is defined similarly, except a sulfur atom, rather than an oxygen atom, is present.

The term "hydroxy" is defined as -OH.

The term "hydroxyalkyl" is defined as a hydroxy group appended to an alkyl group.

The term "amino" is defined as $-\text{NH}_2$, and the term "alkylamino" is defined as $-\text{NR}_2$, wherein at least one R is alkyl and the second R is alkyl or hydrogen.

The term "acylamino" is defined as $\text{RC}(=\text{O})\text{N}$, wherein R is alkyl or aryl.

The term "alkylthio" is defined as -SR, where R is alkyl.

The term "alkylsulfinyl" is defined as R-SO_2 , where R is alkyl.

- 7 -

The term "alkylsulfonyl" is defined as $R-SO_3$, where R is alkyl.

The term "nitro" is defined as $-NO_2$.

5 The term "trifluoromethyl" is defined as $-CF_3$.

The term "trifluoromethoxy" is defined as $-OCF_3$.


The term "cyano" is defined as $-CN$.

10 The term "spiro" as used herein refers to a group having two carbon atoms directly bonded to a carbon atom of the tetracyclic ring system.


In preferred embodiments, R^4 and R^5 are selected from the group consisting of C_{1-6} alkyl, C_{1-6} alkyleneC(=O)OR^a, C(=O)OR^a, C(=S)NR^aR^b, C_{5-7} spiro
15 when combined with R^3 , C_{1-4} alkyleneC(=O)NR^aR^b, C_{1-4} alkyleneNR^aR^b, C_{1-4} alkyleneOR^a, C_{1-4} alkyleneSO₂NR^aR^b, C_{1-4} alkyleneheteroaryl, haloC₁₋₄alkyl, C_{1-4} alkyleneHet, C_{1-4} alkyleneC(=O) C_{1-4} alkyleneHet, and C_{1-4} alkylenearyl, and R^a and R^b, independently, are selected from the
20 group consisting of C_{1-8} alkyl, hydrogen, C_{1-4} alkylenearyl, Het, C_{1-4} alkyleneHet, C_{3-8} cycloalkyl, and haloC₁₋₆alkyl. Het, aryl, and heteroaryl can be substituted with one or more of C(=O)OR^a, NR^aR^b, and C_{1-6} alkyl.

25 In a more preferred group of compounds of formula (I), X is represented by CH₂CH₂, NH, or NCH₃; R³ is hydrogen or methyl; and R⁴ is selected from the group consisting of C_{1-4} alkyl, C_{1-4} alkyleneC(=O)NH₂, C_{1-4} alkyleneNH₂, C_{1-4} alkyleneC(=O)OC₁₋₇alkyl,
30 C_{1-6} alkyleneC(=O)OR^a, C_{1-3} alkyleneOH, C_{1-4} alkyleneSO₂NH₂, C_{1-4} alkyleneOC₁₋₃alkylenearyl, haloC₁₋₃alkyl, C_{1-3} alkylenearyl, C_{1-4} alkyleneC(=O)OC₁₋₃haloalkyl,


- 8 -




5




10




15




20



25



30




- 9 -

In another preferred embodiment, R^3 and R^4 are taken together to form a 5- or 6-membered spiro group.


A preferred R^2 group is represented by

5



10


The bicyclic ring



15

can represent, for example, naphthalene or indene, or a heterocyclic, such as benzoxazole, benzothiazole, benzisoxazole, benzimidazole, quinoline, indole, benzothiophene, or benzofuran, or

20




25

wherein n is an integer 1 or 2, and Y and Z , independently, represent CH_2 , O , S , or NR^a . In a preferred group of compounds of formula (I), R^2 is represented by an optionally substituted bicyclic ring

30

- 10 -




5

wherein n is 1 or 2, and Y and Z are each CH_2 or O.


Especially preferred R^2 bicyclic rings include

10




, or

15



, or

20



25


Within this particular group of compounds, nonlimiting examples of substituents are halogen (e.g., chlorine), hydroxy, C_{1-3} alkyl (e.g., methyl, ethyl, or i-propyl), C_{1-3} alkoxy (e.g., methoxy or ethoxy), CO_2R^b , halomethyl (e.g., trifluoromethoxy), cyano,

30

nitro, and NR^aR^b .

An especially preferred subclass of compounds within the general scope of formula (I) is represented by compounds of formula (II)

- 11 -



(II)

and pharmaceutically acceptable salts and solvates (e.g., hydrates) thereof.

Compounds of formula (I) can contain one or more asymmetric center, and, therefore, can exist as stereoisomers. The present invention includes both mixtures and separate individual stereoisomers of the compounds of formula (I). Compounds of formula (I) also can exist in tautomeric forms, and the invention includes both mixtures and separate individual tautomers thereof.

Pharmaceutically acceptable salts of the compounds of formula (I) can be acid addition salts formed with pharmaceutically acceptable acids. Examples of suitable salts include, but are not limited to, the hydrochloride, hydrobromide, sulfate, bisulfate, phosphate, hydrogen phosphate, acetate, benzoate, succinate, fumarate, maleate, lactate, citrate, tartrate, gluconate, methanesulfonate, benzenesulfonate, and p-toluenesulfonate salts. The compounds of the formula (I) also can provide pharmaceutically acceptable metal salts, in particular alkali metal salts and alkaline earth metal salts, with bases. Examples include the sodium, potassium, magnesium, and calcium salts.

- 12 -

Compounds of the present invention are potent and selective inhibitors of cGMP-specific PDE5. Thus, compounds of formula (I) are of interest for use in therapy, specifically for the treatment of a variety of conditions where selective inhibition of PDE5 is considered to be beneficial.

Phosphodiesterases (PDEs) catalyze the hydrolysis of cyclic nucleotides, such as cyclic adenosine monophosphate (cAMP) and cyclic guanosine monophosphate (cGMP). The PDEs have been classified into at least seven isoenzyme families and are present in many tissues (J.A. Beavo, *Physiol. Rev.*, 75, p. 725 (1995)).

PDE5 inhibition is a particularly attractive target. A potent and selective inhibitor of PDE5 provides vasodilating, relaxing, and diuretic effects, all of which are beneficial in the treatment of various disease states. Research in this area has led to several classes of inhibitors based on the cGMP basic structure (E. Sybertz et al., *Expert. Opin. Ther. Pat.*, 7, p. 631 (1997)).

The biochemical, physiological, and clinical effects of PDE5 inhibitors therefore suggest their utility in a variety of disease states in which modulation of smooth muscle, renal, hemostatic, inflammatory, and/or endocrine function is desirable. The compounds of formula (I), therefore, have utility in the treatment of a number of disorders, including stable, unstable, and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, congestive heart failure, acute respiratory distress syndrome, acute and chronic renal failure, atherosclerosis, conditions of reduced blood vessel

- 13 -

patency (e.g., postpercutaneous transluminal coronary or carotid angioplasty, or post-bypass surgery graft stenosis), peripheral vascular disease, vascular disorders, such as Raynaud's disease, thrombocytopenia, inflammatory diseases, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma, osteoporosis, preterm labor, benign prostatic hypertrophy, peptic ulcer, male erectile dysfunction, female sexual dysfunction, and diseases characterized by disorders of gut motility (e.g., irritable bowel syndrome).

An especially important use is the treatment of male erectile dysfunction, which is one form of impotence and is a common medical problem.

Impotence can be defined as a lack of power, in the male, to copulate, and can involve an inability to achieve penile erection or ejaculation, or both. The incidence of erectile dysfunction increases with age, with about 50% of men over the age of 40 suffering from some degree of erectile dysfunction.

In addition, a further important use is the treatment of female arousal disorder. Female arousal disorders are defined as a recurrent inability to attain or maintain an adequate lubrication/swelling response of sexual excitement until completion of sexual activity. The arousal response consists of vasocongestion in the pelvis, vaginal lubrication, and expansion and swelling of external genitalia.

It is envisioned, therefore, that compounds of formula (I) are useful in the treatment of male erectile dysfunction and female arousal disorder. Thus, the present invention concerns the use

- 14 -

of compounds of formula (I), or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition containing either entity, for the manufacture of a medicament for the curative or prophylactic treatment of erectile dysfunction in a male animal and arousal disorder in a female animal, including humans.

The term "treatment" includes preventing, lowering, stopping, or reversing the progression or severity of the condition or symptoms being treated. As such, the term "treatment" includes both medical therapeutic and/or prophylactic administration, as appropriate.

It also is understood that "a compound of formula (I)," or a physiologically acceptable salt or solvate thereof, can be administered as the neat compound, or as a pharmaceutical composition containing either entity.

Although the compounds of the invention are envisioned primarily for the treatment of sexual dysfunction in humans, such as male erectile dysfunction and female arousal disorder, they also can be used for the treatment of other disease states.

A further aspect of the present invention, therefore, is providing a compound of formula (I) for use in the treatment of stable, unstable, and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, chronic obstructive pulmonary disease, congestive heart failure, acute respiratory distress syndrome, acute and chronic renal failure, atherosclerosis, conditions of reduced blood vessel patency (e.g., post-PTCA or post-bypass graft stenosis), peripheral vascular disease, vascular disorders such

- 15 -

as Raynaud's disease, thrombocythemia, inflammatory diseases, prophylaxis of myocardial infarction, prophylaxis of stroke, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma, osteoporosis, preterm labor, benign prostatic hypertrophy, male and female erectile dysfunction, or diseases characterized by disorders of gut motility (e.g., IBS).

According to another aspect of the present invention, there is provided the use of a compound of formula (I) for the manufacture of a medicament for the treatment of the above-noted conditions and disorders.

In a further aspect, the present invention provides a method of treating the above-noted conditions and disorders in a human or nonhuman animal body which comprises administering to said body a therapeutically effective amount of a compound of formula (I).

Compounds of the invention can be administered by any suitable route, for example by oral, buccal, inhalation, sublingual, rectal, vaginal, transurethral, nasal, topical, percutaneous, i.e., transdermal, or parenteral (including intravenous, intramuscular, subcutaneous, and intracoronary) administration. Parenteral administration can be accomplished using a needle and syringe, or using a high pressure technique, like POWDERJECT™.

Oral administration of a compound of the invention is the preferred route. Oral administration is the most convenient and avoids the disadvantages associated with other routes of administration. For patients suffering from a swallowing dis-

- 16 -

order or from impairment of drug absorption after oral administration, the drug can be administered parenterally, e.g., sublingually or buccally.

Compounds and pharmaceutical compositions suitable for use in the present invention include those wherein the active ingredient is administered in an effective amount to achieve its intended purpose. More specifically, a "therapeutically effective amount" means an amount effective to prevent development of, or to alleviate the existing symptoms of, the subject being treated. Determination of the effective amounts is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

A "therapeutically effective dose" refers to that amount of the compound that results in achieving the desired effect. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index, which is expressed as the ratio between LD₅₀ and ED₅₀. Compounds which exhibit high therapeutic indices are preferred. The data obtained from such data can be used in formulating a dosage range for use in humans. The dosage of such compounds preferably lies within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage can vary within this range

- 17 -

depending upon the dosage form employed, and the route of administration utilized.

5 The exact formulation, route of administration, and dosage can be chosen by the individual physician in view of the patient's condition. Dosage amount and interval can be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the therapeutic effects.

10 The amount of composition administered is dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration, and the judgment of the prescribing physician.

15 Specifically, for administration to a human in the curative or prophylactic treatment of the conditions and disorders identified above, oral dosages of a compound of formula (I) generally are about 0.5 to about 1000 mg daily for an average adult patient (70 kg). Thus, for a typical adult
20 patient, individual tablets or capsules contain 0.2 to 500 mg of active compound, in a suitable pharmaceutically acceptable vehicle or carrier, for administration in single or multiple doses, once or several times per day. Dosages for intravenous,
25 buccal, or sublingual administration typically are 0.1 to 500 mg per single dose as required. In practice, the physician determines the actual dosing regimen which is most suitable for an individual patient, and the dosage varies with the age, weight,
30 and response of the particular patient. The above dosages are exemplary of the average case, but there can be individual instances in which higher or lower

- 18 -

dosages are merited, and such are within the scope of this invention.

For human use, a compound of the formula (I) can be administered alone, but generally is administered in admixture with a pharmaceutical carrier selected with regard to the intended route of administration and standard pharmaceutical practice. Pharmaceutical compositions for use in accordance with the present invention thus can be formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries that facilitate processing of compounds of formula (I) into preparations which can be used pharmaceutically.

These pharmaceutical compositions can be manufactured in a conventional manner, e.g., by conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes. Proper formulation is dependent upon the route of administration chosen. When a therapeutically effective amount of a compound of the present invention is administered orally, the composition typically is in the form of a tablet, capsule, powder, solution, or elixir. When administered in tablet form, the composition can additionally contain a solid carrier, such as a gelatin or an adjuvant. The tablet, capsule, and powder contain about 5 to about 95% compound of the present invention, and preferably from about 25 to about 90% compound of the present invention. When administered in liquid form, a liquid carrier such as water, petroleum, or oils of animal or plant origin can be added. The liquid form of

- 19 -

the composition can further contain physiological saline solution, dextrose or other saccharide solutions, or glycols. When administered in liquid form, the composition contains about 0.5 to about 5 90% by weight of a compound of the present invention, and preferably about 1 to about 50% of a compound of the present invention.

When a therapeutically effective amount of a compound of the present invention is administered 10 by intravenous, cutaneous, or subcutaneous injection, the composition is in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. 15 A preferred composition for intravenous, cutaneous, or subcutaneous injection typically contains, in addition to a compound of the present invention, an isotonic vehicle.

20 For oral administration, the compounds can be formulated readily by combining a compound of formula (I) with pharmaceutically acceptable carriers well known in the art. Such carriers enable the present compounds to be formulated as 25 tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by adding a compound of formula (I) with a solid excipient, optionally grinding a resulting mixture, and 30 processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients include, for

- 20 -

example, fillers and cellulose preparations. If desired, disintegrating agents can be added.

For administration by inhalation, compounds of the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant. In the case of a pressurized aerosol, the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g., gelatin, for use in an inhaler or insufflation can be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds can be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection can be presented in unit dosage form, e.g., in ampules or in multidose containers, with an added preservative. The compositions can take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and can contain formulatory agents such as suspending, stabilizing, and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds can be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils or synthetic fatty acid esters. Aqueous injection suspensions can contain substances which increase the viscosity of the suspension. Optionally, the suspension also can contain suitable

- 21 -

stabilizers or agents that increase the solubility of the compounds and allow for the preparation of highly concentrated solutions. Alternatively, a present composition can be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

Compounds of the present invention also can be formulated in rectal compositions, such as suppositories or retention enemas, e.g., containing conventional suppository bases. In addition to the formulations described previously, the compounds also can be formulated as a depot preparation. Such long-acting formulations can be administered by implantation (for example, subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds can be formulated with suitable polymeric or hydrophobic materials (for example, as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

Many of the compounds of the present invention can be provided as salts with pharmaceutically compatible counterions. Such pharmaceutically acceptable base addition salts are those salts that retain the biological effectiveness and properties of the free acids, and that are obtained by reaction with suitable inorganic or organic bases.

In particular, a compound of formula (I) can be administered orally, buccally, or sublingually in the form of tablets containing excipients, such as starch or lactose, or in capsules or ovules, either alone or in admixture with excipients, or in

- 22 -

the form of elixirs or suspensions containing
flavoring or coloring agents. Such liquid prepara-
tions can be prepared with pharmaceutically accept-
able additives, such as suspending agents. A com-
5 compound also can be injected parenterally, for
example, intravenously, intramuscularly, subcutane-
ously, or intracoronarily. For parenteral adminis-
tration, the compound is best used in the form of a
sterile aqueous solution which can contain other
10 substances, for example, salts, or monosaccharides,
such as mannitol or glucose, to make the solution
isotonic with blood.

For veterinary use, a compound of formula
(I) or a nontoxic salt thereof, is administered as a
15 suitably acceptable formulation in accordance with
normal veterinary practice. The veterinarian can
readily determine the dosing regimen and route of
administration that is most appropriate for a
particular animal.


20 Thus, the invention provides in a further
aspect a pharmaceutical composition comprising a
compound of the formula (I), together with a pharma-
ceutically acceptable diluent or carrier therefor.
There is further provided by the present invention a
25 process of preparing a pharmaceutical composition
comprising a compound of formula (I), which process
comprises mixing a compound of formula (I), together
with a pharmaceutically acceptable diluent or
carrier therefor.

30 In a particular embodiment, the invention
includes a pharmaceutical composition for the cura-
tive or prophylactic treatment of erectile dysfunc-
tion in a male animal, or arousal disorder in a

- 23 -

female animal, including humans, comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable diluent or carrier.


Compounds of formula (I) can be prepared by any suitable method known in the art, or by the following processes which form part of the present invention. In the methods below, R^0 through R^5 , R^a , R^b , and X are as defined in structural formula (I) above. In particular, Daugan U.S. Patent No. 5,859,006, incorporated herein by reference, discloses preparation of a compound of structural formula (III).



(III)


In short, the compound of structural formula (III), i.e., the *cis*-isomer of Intermediates 1 and 2 of Daugan U.S. Patent No. 5,859,006 was prepared according to the following reaction scheme:

- 24 -




5

D-Tryptophan methyl ester




10

Piperonal



15

20




25

30

(42%)

(III) (cis-isomer)

- 25 -



(28%)

(trans-isomer)

A compound of structural formula (I) is prepared similarly by reacting a tryptophan ester, or a tryptophan ester substituted with suitable R^0 substituents, with a suitable aldehyde to provide the desired R^2 substituent. The resulting product then is cyclized by reaction with a suitable amine to provide a compound of structural formula (I). The cyclization reaction is disclosed in Daugan U.S. Patent No. 5,859,006.

In the synthesis of compounds of structural formula (I), protecting compounds and protecting groups, like benzyl chloroformate and trichloroethyl chloroformate, which are well known to persons skilled in the art, can be used. Such protecting groups are disclosed, for example, in T.W. Greene et al. "Protective Groups in Organic Synthesis, Third Edition," John Wiley and Sons, Inc., NY, NY (1999). These protecting groups are removed in the final steps of the synthesis under basic, acidic, or hydrogenolytic conditions which are readily apparent to those skilled in the art. By employing appropri-

- 26 -

ate manipulation and protection of chemical functionalities, synthesis of compounds of structural formula (I) not specifically set forth herein can be accomplished by methods analogous to the schemes set forth below. For example, the structure of a compound of structural formula (I) can be varied by using an appropriate aldehyde to change the identity of R², or by using a halo or alkyl phenyl-substituted tryptophan ester.

Compounds of formula (I) can be converted to other compounds of formula (I). Thus, for example, when a compound contains a substituted aromatic ring, it is possible to prepare another suitably substituted compound of formula (I).

Examples of appropriate interconversions include, but are not limited to, OR^a to hydroxy by suitable means (e.g., using a agent such as SnCl₂ or a palladium catalyst, such as palladium-on-carbon), or amino to substituted amino, such as alkylamine, using standard acylating or sulfonylating conditions.



Compounds of structural formula (I) can be made accordingly to one of the following four non-limiting synthetic routes.

Method A


The 1,2,3,4-tetrahydro- β -carboline of general formula (IV) can be prepared by the Pictet-Spengler reaction as set forth in Daugan U.S. Patent No. 5,859,006 and in A. Madrigal et al., *J. Org. Chem.*, 63, page 2724 (1998), for example. The resulting secondary amine then is treated with either an amino acid or an acid halide, under

- 27 -

suitable acylation conditions, to form an amide-
ester. Ring cyclization to form the diketo-
piperazine (I) is accomplished by intramolecular
amine attack on the ester. The amine can be derived
5 from a suitable side chain bearing a leaving group
that reacts with primary amine compound (V).




- 28 -



5

or




10

Method B

The diketopiperazine (I) also can be prepared by reaction of a tryptophan with an amino acid under typical peptide coupling conditions to form an N-acyltryptophan. Ring cyclization to form diketopiperazine (I) is accomplished by intramolecular amine attack on the ester. The resulting piperazine can undergo a condensation with an aldehyde under modified Pictet-Spengler conditions (see T.A. Miller et al., *Bioorg. Med. Chem. Lett.*, 8, p. 1065 (1998), for example) to give the β -carboline.

- 29 -

**Method C**

5

The β -carboline skeleton also can be prepared using the Bischler-Napieralski reaction, which is a cyclodehydration of an acylated tryptophan, as disclosed in W.M. Whaley et al., *Org. React.*, VI, pp. 74-150 (1951). P_2O_5 and POCl_3 are the most common cyclization reagents. Reduction of the imine with NaBH_4 , for example, yields the 1,2,3,4-tetrahydro- β -carboline.



10

15


A modified method, which avoids the potential for racemisation, includes acylating the amine

- 30 -


of tryptophan first, followed by conversion to the thioamide with, for example, with Lawesson's reagent. Treatment of the thioamide with an alkyl halide or acyl halide provides the iminium halide. Reduction of the crude iminium halide with sodium borohydride (NaBH_4) at reduced temperature stereoselectively provides the 1,2,3,4-tetrahydro- β -carboline.



- 31 -




10



or


15



Lg=leaving group

20



25




30

- 32 -

5




10



or

15




Lg=leaving group

20

- 33 -

Method D

In addition, the β -carboline can be treated with a chloroformate to form a 2,3-dicarboxylic acid diester. Treatment of the diester with, for example, a hydrazine, provides compounds of general formula (I) when X contains nitrogen (see K. Winterfield et al., *Arch. Pharmaz*, 304, 216 (1971) and E.H. White et al., *J. Org. Chem.*, 32, 1921 (1967)).



Compounds of formula (I) can be prepared by the methods above as individual stereoisomers from the appropriate stereoisomer of formula (III) or as a racemic mixture from the appropriate racemic compound of formula (III). Individual stereoisomers of the compounds of the invention can be prepared

- 34 -

from racemates by resolution using methods known in the art for the separation of racemic mixtures into their constituent stereoisomers, for example, using HPLC on a chiral column, such as Hypersil naphthyl urea, or using separation of salts of stereoisomers. Compounds of the invention can be isolated in association with solvent molecules by crystallization from, or evaporation of, an appropriate solvent.

The pharmaceutically acceptable acid addition salts of the compounds of formula (I) that contain a basic center can be prepared in a conventional manner. For example, a solution of the free base can be treated with a suitable acid, either neat or in a suitable solution, and the resulting salt isolated either by filtration or by evaporation under vacuum of the reaction solvent. Pharmaceutically acceptable base addition salts can be obtained in an analogous manner by treating a solution of a compound of formula (I) with a suitable base. Both types of salt can be formed or interconverted using ion-exchange resin techniques. Thus, according to a further aspect of the invention, a method for preparing a compound of formula (I) or a salt or solvate (e.g., hydrate) is provided, followed by (i) salt formation, or (ii) solvate (e.g., hydrate) formation.


The following abbreviations are used hereafter in the accompanying examples: rt (room temperature), min (minute), h (hour), g (gram), mmol (millimole), m.p. (melting point), eq (equivalents), quant (quantitative), L (liter), mL (milliliter), μ L (microliters), DCC (dicyclohexylcarbodiimide), Cbz

- 35 -

(benzylcarbamate), DMSO (dimethyl sulfoxide), CH_2Cl_2 (dichloromethane), CHCl_3 (chloroform), IPA (isopropyl alcohol), TFA (trifluoroacetic acid), EtOH (ethanol), AcOH (acetic acid), EtOAc (ethyl acetate), MeOH (methanol), Et_3N (triethylamine), DMF (dimethylformamide), EtOAc (ethyl acetate), and THF (tetrahydrofuran).

The following illustrates specific examples of compounds of structural formula (I) and synthetic routes to some of these structures.

Preparation of Example 1




Example 1


Example 1 was prepared from a tryptophan ester (Intermediate 1). The tryptophan ester utilized to prepare Example 1 is available commercially from Aldrich Chemical Co., Milwaukee, WI. The Pictet-Spengler reaction to provide Example 1 from Intermediate 4 is disclosed in T.A. Miller et al., *Bioorg. Med. Chem. Lett.*, 8, 1065 (1998); J.J. Tepe et al., *J. Med. Chem.*, 39, 2188 (1996); C.W. Ong et al., *Aust. J. Chem.*, 43, 773 (1990); and R.H. Herbert et al., *Tetrahedron Lett.*, 35, 5497 (1994).

- 36 -

5




10




15

Intermediate 2



Intermediate 3


20



25

Intermediate 4

- 37 -



Example 1
(1:1 Mixture of Diastereomers)

Preparation of Cbz-Protected
Amide Intermediate 2

Dicyclohexylcarbodiimide (1.8 g, 8.8 mmol) was added to a solution of N-Cbz-N,2-dimethylalanine (2.0 g, 8.0 mmol) in methylene chloride (10 mL) at 0°C under a nitrogen blanket in one portion. The resulting white slurry was stirred at 0°C for 30 min, after which a slurry of Intermediate 1 (1.7 g, 8.0 mmol) and triethylamine (1.3 mL, 10 mmol) in methylene chloride (10 mL) was added over 5 min. The resulting white slurry was slowly warmed to room temperature, then stirred for a total of 6 days. The slurry then was filtered under reduced pressure, and the solid was washed with CH₂Cl₂ (50 mL). The filtrate was concentrated under reduced pressure to provide a yellow oil, which was purified by flash column chromatography, eluting with ethyl acetate/-chloroform (1:9), to provide Cbz-protected amide

- 38 -

Intermediate 2 as a white semi-solid (3.76 g, 100%):
TLC R_f (1:4 ethyl acetate/chloroform)=0.35; ^1H NMR
(300 MHz, $\text{DMSO}-d_6$): δ 10.85 (s, 1H), 7.76 (d, $J=7.6$
Hz, 1H), 7.46 (d, $J=7.6$ Hz, 1H), 7.34-7.26 (m, 5H),
5 7.12 (s, 1H), 7.06-6.94 (m, 2H), 4.88-4.83 (m, 1H),
4.46 (q, $J=8.0$ Hz, 1H), 3.53 (s, 3H), 3.09 (t, $J=6.5$
Hz, 2H), 2.89 (t, 3H), 1.29 (s, 3H), 1.23 (s, 3H)
ppm.

10 Preparation of Methylamine Intermediate 3

A solution of Intermediate 2 (3.7 g, 8.2
mmol) in acetic acid (10 mL) and methanol (50 mL)
was treated with a catalytic amount of 10% palladium
15 on carbon (0.40 g, 50% wet) and the resulting mix-
ture was stirred under a hydrogen atmosphere at 50°C
for 29 hours. The reaction mixture then was cooled
to room temperature, and the palladium catalyst was
removed by vacuum filtration through a plug of
20 Celite. The filtrate was concentrated under reduced
pressure to provide a yellow residue, which was
purified by flash column chromatography, eluting
with methanol/chloroform (1:9), to provide Inter-
mediate 3 as an amber oil (1.65 g, 65%): TLC R_f (1:9
25 methanol/chloroform)=0.55; ^1H NMR (300 MHz, CDCl_3):
 δ 8.17 (s, 1H), 7.70 (d, $J=7.9$ Hz, 1H), 7.59 (d,
 $J=7.8$ Hz, 2H), 7.35 (d, $J=7.7$ Hz, 1H), 7.19-7.12 (m,
2H), 4.87 (q, $J=5.3$ Hz, 1H), 3.72 (s, 3H), 3.37-3.30
(m, 2H), 2.27 (s, 3H), 1.38 (s, 3H), 1.33 (s, 3H)
30 ppm; API MS m/z 318 $[\text{C}_{17}\text{H}_{23}\text{N}_3\text{O}_3+\text{H}]^+$.

- 39 -

Preparation of Diketopiperazine Intermediate 4

A suspension of Intermediate 3 (1.6 g, 5.0 mmol) in ethanol (50 mL) was heated to reflux under a nitrogen blanket and stirred for a total of 3 days. The solvent was removed under reduced pressure to provide a residue which was purified by flash column chromatography, eluting with methanol/chloroform (1:19), to provide diketopiperazine Intermediate 4 as a white foam (0.810 g, 56%): mp 223-229°C; TLC R_f (1:9 methanol/chloroform)=0.63; ^1H NMR (300 MHz, DMSO- d_6): δ 10.86 (s, 1H), 8.12 (s, 1H), 7.50 (d, $J=7.9$ Hz, 1H), 7.28 (d, $J=8.0$ Hz, 1H), 7.01-6.93 (m, 3H), 4.19 (s, 1H), 3.29-3.28 (m, 1H), 2.98 (dd, $J=4.6, 14.2$ Hz, 1H), 2.57 (s, 3H), 1.22 (s, 6H) ppm; CI MS (methane) m/z 286 [$\text{C}_{16}\text{H}_{19}\text{N}_3\text{O}_2+\text{H}$] $^+$; $[\alpha]_D^{25^\circ\text{C}}=-26.5^\circ$ ($c=0.10$, DMSO).

Preparation of Example 1


(12a,R)-6-Benzo[1,3]dioxol-5-yl,2,3,3-trimethyl-2,3,6,7,12,12a-hexahydropyrazino[1',2':1,6]-pyrido[3,4-b]indole-1,4-dione

Trifluoroacetic acid (0.5 mL, 6.2 mmol) was added to a suspension of Intermediate 4 (0.800 g, 2.8 mmol) and piperonal (0.505 g, 3.4 mmol) in methylene chloride (10 mL) at 0°C under a nitrogen blanket, after which the solution was warmed to room temperature and stirred for a total of 22 hours. The reaction mixture was diluted with ethyl acetate (150 mL), washed with saturated sodium bicarbonate (NaHCO_3) solution (100 mL) and brine (50 mL), treated with silica gel (5 g), then the solvent was removed

- 40 -



under reduced pressure. The residue was purified by flash column chromatography, eluting with methanol/chloroform (1:49), to provide Example 1 as a white powder (0.534 g, 46%, 1:1 mixture of *cis* and *trans* diastereomers at C6): mp 225-235°C; TLC R_f (1:19 methanol/chloroform)=0.87; ^1H NMR (300 MHz, DMSO- d_6): δ 10.08 (s, 2H), 8.37 (s, 1H), 8.25 (s, 1H), 7.46 (t, $J=8.9$ Hz, 2H), 7.28 (t, $J=8.2$ Hz, 2H), 7.02-6.95 (m, 4H), 6.80 (d, $J=8.0$ Hz, 2H), 6.60 (s, 1H), 6.48 (d, $J=8.1$ Hz, 1H), 6.15 (s, 1H), 5.98 (d, $J=4.4$ Hz, 2H), 4.19 (s, 2H), 3.22 (dd, $J=3.9, 14.8$ Hz, 2H), 3.00-2.89 (m, 2H), 2.70 (s, 3H), 2.62 (s, 3H), 1.35 (s, 3H), 1.31 (s, 3H), 0.65 (s, 3H), 0.47 (s, 3H) ppm; API MS m/z 418 [$\text{C}_{24}\text{H}_{23}\text{N}_3\text{O}_4+\text{H}$] $^+$; $[\alpha]_D^{25^\circ\text{C}}=-63.1^\circ$ ($c=0.50$, CHCl_3). Anal. Calcd. for $\text{C}_{24}\text{H}_{23}\text{N}_3\text{O}_4$: C, 69.05; H, 5.55; N, 10.07. Found: C, 68.91; H, 5.56; N, 10.04. Chiral HPLC analysis (Chiralcel OD Column, 250 x 4.6 mm, Retention Time=7.34 min; 1:1 isopropanol/hexanes; flow=0.5 mL/min; detector @ 254 nm; 25°C) showed one major peak, with a purity of 99.9%.

Preparation of Example 2




- 41 -

Example 2 was prepared from the compound of structural formula (III) and N-Cbz-L-glutamine as illustrated below.



Intermediate 6



- 42 -

**Preparation of *cis*- β -Carboline
Amide Intermediate 6**

5 To a solution of N-Cbz-L-glutamine
(Intermediate 5) (5.0 g, 17.8 mmol) in THF (80 mL)
at 0°C under a nitrogen blanket was added dicyclo-
hexylcarbodiimide (4.4 g, 21.4 mmol) in one portion.
The resulting white slurry was stirred at 0°C for 30
10 minutes, after which Compound (III) (6.3 g, 17.8
mmol) was added in one portion. The resulting amber
slurry was slowly warmed to room temperature, then
stirred 48 hours. The slurry was filtered under re-
duced pressure and the solid was washed with methyl-
15 ene chloride (500 mL). The filtrate was concen-
trated under reduced pressure to provide an amber
foam, which was partially purified by flash column
chromatography, eluting with methanol/chloroform
(1:9), to provide Intermediate 6 as an off-white
20 foam (4.6 g): TLC R_f (9:1 methanol/chloroform)=0.60.

Preparation of Example 2


3-((3S,6R,12aR)-6-Benzo[1,3]dioxol-5-yl-1,4-dioxo-
1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]-
25 pyrido[3,4-b]indo-3-yl)propionamide

A solution of Intermediate 6 (4.6 g, 7.5
mmol) in methanol (50 mL) and acetic acid (5 mL) was
30 treated with a catalytic amount of 10% palladium on
carbon (500 mg, 50% wet) and the resulting mixture
was stirred under a hydrogen atmosphere at 50°C for
3 hours. The reaction mixture then was cooled to
room temperature and the palladium catalyst was re-
35 moved by vacuum filtration through Celite (10 g).
The filtrate was concentrated under reduced pressure

- 43 -

and the residue was purified by flash column chromatography, eluting with methanol/chloroform (1:9), to provide Example 2 as a white powder (1.24 g, 15% over two steps): mp 78-82°C; TLC R_f (4:1 chloroform/methanol)=0.58; ^1H NMR (300 MHz, $\text{DMSO}-d_6$): δ 11.05 (s, 1H), 8.57 (d, $J=3.8$ Hz, 1H), 7.52 (d, $J=7.3$ Hz, 1H), 7.30 (d, $J=7.7$ Hz, 1H), 7.23 (s, 1H), 7.08-6.97 (m, 2H), 6.78 (s, 1H), 6.10 (s, 1H), 5.92 (s, 2H), 4.43 (dd, $J=4.4, 11.4$ Hz, 1H), 3.83-3.80 (m, 1H), 3.61 (dd, $J=4.5, 15.8$ Hz, 1H), 3.32 (s, 3H), 2.93 (dd, $J=11.7, 15.8$ Hz, 1H), 2.17 (t, $J=7.2$ Hz, 2H), 1.94 (t, $J=7.4$ Hz, 2H) ppm; API MS m/z 447 $[\text{C}_{24}\text{H}_{22}\text{N}_4\text{O}_5+\text{H}]^+$. Anal. Calcd. for $\text{C}_{24}\text{H}_{22}\text{N}_4\text{O}_5\cdot\text{H}_2\text{O}$: C, 62.06; H, 5.21; N, 12.06. Found: C, 62.12; H, 5.00; N, 12.07. The relative stereochemistry of analog Example 2 was confirmed to be the *cis* isomer by a series of NOE difference experiments: positive NOE enhancements from the C12a proton at 4.45 ppm to the C6 proton at 6.10 ppm; a positive NOE enhancement from the C6 proton at 6.10 ppm to the C12a proton at 4.45 ppm.



Preparation of Example 3



- 44 -


Example 3 was prepared from the hydrochloride of Compound (III) by the following synthetic sequence. The 1,2,4-triazine-3,6-dione of Example 3 was prepared by the method of K.

5 Winterfield et al., *Arch. Pharmaz.*, 304, 0. 216 (1971).



Intermediate 7

- 45 -



Example 3

10

15

**Preparation of *cis*- β -Carboline
Carbamate Intermediate 7**

15

20

25

Methyl chloroformate (4.8 mL, 62 mmol) was added dropwise to a suspension of Compound (III) (20 g, 52 mmol) and N-methylmorpholine (14.2 mL, 129 mmol) in THF (150 mL) at 0°C under a nitrogen blanket. The mixture was slowly warmed to room temperature, then stirred for 3 days. The resulting mixture was diluted with ethyl acetate (200 mL), washed with brine (150 mL), dried over magnesium sulfate (MgSO_4), and filtered. The solvent was removed under reduced pressure to afford Intermediate 7 as an amber foam (21 g, 96%), which was suitable for use without further purification.

- 46 -

Preparation of Example 3

(5aR,10R)-10-Benzo[1,3]dioxol-5-yl-5,5a,7,8,10,11-hexahydro-7,8,9a,11-tetraazabenzob[b]fluorene-6,9-dione

5

Sodium methoxide (5.2 mL, 27 mmol, 30% solution in methanol) was added dropwise to a mixture of Intermediate 7 (5 g, 12 mmol) and anhydrous hydrazine (0.5 mL, 15 mmol) in ethanol (30 mL), and the mixture was heated at reflux under a nitrogen blanket for 27 hours. The suspension was cooled to room temperature and the orange solids were removed by vacuum filtration. The organic phase was concentrated under reduced pressure to afford an orange foam which was purified by reverse phase flash column chromatography on C-18 silica gel, eluting with methanol/water (1:2), to provide the crude product as an orange oil. The oil was further purified by flash column chromatography, eluting with methanol/chloroform (1:4), to provide Example 3 as a yellow powder (0.465 g, 10%): mp 188-194°C; TLC R_f (1:4 methanol/chloroform)=0.37; ¹H NMR (300 MHz, D₂O): δ 7.72 (d, J=7.6 Hz, 1H), 7.42 (t, J=8.6 Hz, 1H), 7.30-7.23 (m, 2H), 7.00 (s, 1H), 6.23-6.83 (m, 1H), 6.04 (d, J=11.1 Hz, 2H), 5.73 (s, 1H), 4.18 (dd, J=7.0, 11.5 Hz, 1H), 3.51 (dd, J=5.0, 11.4 Hz, 1H), 3.22 (t, J=15.3 Hz, 1H) ppm; API MS m/z 377 [C₂₀H₁₆N₄O₄+H]⁺; [α]_D^{25°C}=+44.7° (c=0.5, methanol).

Anal. Calcd. for C₂₀H₁₆N₄O₄·0.25 H₂O: C, 63.07; H, 4.37; N, 14.71. Found: C, 63.06; H, 4.30; N, 14.71. The relative stereochemistry of Example 3 was confirmed to be the *cis* isomer by a series of NOE difference experiments: a positive NOE enhance-

30


- 47 -

ment from the C12a proton at 3.57 ppm to the C6 proton at 5.73 ppm; a positive NOE enhancement from the C6 proton at 5.73 ppm to the C12a proton at 3.57 ppm.

5


Preparation of Example 4

10



15


The compound of Example 4 was prepared from Compound (III) by the following reaction scheme.



Intermediate 8

20

- 48 -



Example 4

15

Preparation of (+)-*cis*-3-Chloropropionyl- β -carboline Intermediate 8

20

To a mixture of Compound (III) (6.00 g, 15.5 mmol) and triethylamine (6.5 mL, 46.5 mmol) in THF (100 mL) and water (25 mL), was added 3-chloropropionyl chloride (1.9 mL, 20.2 mmol), dropwise, at 0°C under a nitrogen blanket. The resulting mixture was warmed to room temperature and stirred for 1.5 hours. The reaction was quenched with 1 N HCl (40 mL) and was concentrated to remove THF. The residue

25

was diluted with EtOAc (200 mL), then the layers were separated. The organic phase was washed with 1 N HCl (50 mL), saturated NaHCO₃ (50 mL), and saturated sodium chloride (NaCl) (50 mL), then dried over anhydrous Na₂SO₄. Filtration and concentration *in*

30

vacuo gave an oil which was purified by column chromatography (silica gel, 0-5% EtOAc/CH₂Cl₂). The product was obtained as a pale yellow foam 2.30 g (33.6%): TLC R_f (90:10:1/CH₂Cl₂:EtOAc:MeOH)=0.80; ¹H NMR (300 MHz, CDCl₃): δ 10.86 (s, 1H), 7.54 (d,

- 49 -

J=7.6 Hz, 1H), 7.28 (d, J=7.9 Hz, 1H), 7.10 (t, J=7.4 Hz, 1H), 7.02 (t, J=7.3 Hz, 1H), 6.85 (s, 1H), 6.79 (d, J=8.1 Hz, 1H), 6.67 (s, 1H), 6.44 (d, J=8.0 Hz, 1H), 6.27 (s, 1H), 5.97 (d, J=4.5 Hz, 2H), 5.28 (d, J=6.6 Hz, 1H), 3.90 (t, J=6.5 Hz, 2H), 3.43 (d, J=16.0 Hz, 1H), 3.17-3.37 (m, 2H), 2.91-3.06 (m, 4H).

Preparation of Example 4


10 (5aS,11R)-11-Benzo[1,3]dioxol-5-yl-7-methyl-5,5a,8,9,11,12-hexahydro-7H-7,10a,12-triazacyclohepta[b]fluorene-6,10-dione

15 A mixture of Intermediate 8 (1.22 g, 2.77 mmol), 40% methylamine in water (1.2 mL, 13.8 mmol), and THF (20 mL) was stirred at 45°C under a nitrogen blanket. After 4 hours, only starting material and the 3-methylamino intermediate were detected. Sodium iodide (20.7 mg, 0.138 mmol) and 40% methylamine
20 in water (1.2 mL) were added, and the mixture was stirred at 45°C for 4 days. The mixture was cooled to room temperature and was quenched with 1 N HCl (2 mL). The residue was diluted with CH₂Cl₂ (100 mL),
25 washed with 0.1 N HCl (10 mL), water (30 mL), and saturated NaCl (20 mL), then dried over anhydrous Na₂SO₄. Filtration and concentration *in vacuo* gave a yellow foam which was purified by column chromatography (silica gel, 90:10:1/CH₂Cl₂:EtOAc:MeOH). The
30 product was obtained as a white solid 0.55 g (49.5%) after recrystallization from MeOH/water. The product was contaminated with approximately 8% of the *cis* isomer: mp 179-182°C; TLC R_f (90:10:1/-CH₂Cl₂:EtOAc:MeOH)=0.27; ¹H NMR (300 MHz, DMSO-d₆): δ

- 50 -



11.08 (s, 1H), 7.48 (d, J=7.7 Hz, 1H), 7.32 (d, J=7.9 Hz, 1H), 7.10 (t, J=7.5 Hz, 1H), 7.01 (t, J=7.2 Hz, 1H), 6.86 (d, J=8.0 Hz, 1H), 6.83 (s, 1H), 6.79 (d, J=1.5 Hz, 1H), 6.59 (d, J=8.7 Hz, 1H), 6.01 (d, J=2.3 Hz, 2H), 4.45 (dd, J=4.1 Hz, J=11.0 Hz, 1H), 3.74-3.82 (m, 1H), 3.27-3.45 (m, 2H), 3.08 (dd, J=4.1 Hz, J=15.4 Hz, 1H), 2.91-3.00 (m, 4H), 2.74-2.81 (m, 1H); MS (API) m/z 404 (M+H), 426 (M+Na); $[\alpha]_D^{25} = -140.7^\circ$ (c=0.17, DMSO). Anal. Calcd. for $C_{23}H_{21}N_3O_4 \cdot 0.4 H_2O$: C, 67.27; H, 5.35; N, 10.23. Found: C, 67.28; H, 5.30; N, 10.09. The relative stereochemistry of the major product was confirmed to be the *trans* isomer by NOE difference experiments (DMSO- d_6): no NOE enhancements from the C5a proton at 1.29 ppm to the C11 proton at 6.83 ppm.

Preparation of Example 5




The compound of Example 5 was prepared from Compound (III) and (L)-N-Cbz- γ -*tert*-butyl-glutamic acid by the following synthetic scheme.

- 51 -



Intermediate 9



Example 5

Preparation of Intermediate 9

15

To a solution of L-Cbz- γ -*tert*-butyl-glutamic acid (5.0 g, 14.8 mmol) in methylene chloride (30 mL) at 0°C was added dicyclohexylcarbodiimide

- 52 -

(3.09 g, 15 mmol) in one portion. The resulting white slurry was stirred at 0°C for 30 minutes, after which Compound (III) was added. The resulting yellow slurry was slowly warmed to room temperature, and stirred for a total of 18 hours. The slurry was filtered under reduced pressure and the solid was washed with methylene chloride (5 x 10 mL). The filtrate was concentrated under reduced pressure and the residue was purified by flash column chromatography, eluting with methylene chloride/ethyl acetate (3:1), to provide the impure Intermediate 9 (containing a small amount of dicyclohexylcarbodiimide) as a yellow solid which was used without further purification (8.3 g): TLC R_f (4:1 methylene chloride/ethyl acetate)=0.91.

Preparation of Example 5

3-((3S,6R,12aR)-6-Benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]-pyrido[3,4-b]indol-3-yl-propionic acid tert-butyl ester

Intermediate 9 was dissolved in methanol (100 mL) and treated with a catalytic amount of 5% palladium on carbon (0.50 g, about 50% wet). The mixture was stirred under a hydrogen atmosphere at 50°C for 6 hours, after which the palladium catalyst was removed by vacuum filtration through a plug of Celite, eluting with methanol (100 mL). The filtrate was concentrated under reduced pressure and the residue was purified by flash column chromatography, eluting with methylene chloride/ethyl acetate (4:1), to provide Example 5 as a white solid (5.1 g, 69% over two steps). A small lot of this product


- 53 -

was further purified by recrystallization from methylene chloride at -10°C : mp $222-227^{\circ}\text{C}$; TLC R_f (4:1 methylene chloride/ethyl acetate)=0.40; ^1H NMR (300 MHz, CDCl_3): δ 7.74 (bs, 1H), 7.65-7.56 (m, 1H), 7.30-7.10 (m, 3H), 6.90-6.83 (m, 1H), 6.75-6.67 (m, 2H), 6.26 (d, $J=3.3$ Hz, 1H); 5.19 (s, 1H), 5.89 (s, 1H), 5.86 (s, 1H), 4.41 (dd, $J=5.7, 4.1$ Hz, 1H), 4.10-4.02 (m, 1H), 3.76 (dd, $J=8.0, 4.1$ Hz, 1H), 3.30-3.17 (m, 1H), 2.42 (t, $J=7.1$ Hz, 2H), 2.28-2.00 (m, 2H), 1.44 (s, 9H); ^{13}C NMR (125 MHz, CDCl_3): δ 171.8, 168.7, 167.9, 147.9, 136.5, 135.6, 132.9, 126.3, 122.6, 120.6, 120.2, 118.6, 111.2, 108.3, 107.3, 106.3, 101.2, 81.2, 57.2, 55.4, 55.2, 31.3, 29.4, 28.1, 28.8 ppm; API MS m/z 504 $[\text{C}_{28}\text{H}_{29}\text{N}_3\text{O}_6+\text{H}]^+$; $[\alpha]_D^{25^{\circ}\text{C}}=+31.3^{\circ}$ ($c=1.0$, CHCl_3). Anal. Calcd. for $\text{C}_{28}\text{H}_{29}\text{N}_3\text{O}_6$: C, 66.79; H, 5.80; N, 8.34. Found: C, 66.27; H, 5.74; N, 8.30. The relative stereochemistry of Example 5 was confirmed to be the *cis* isomer by an NOE difference experiment (DMSO- d_6): positive NOE enhancements from the C12a proton at 4.40 ppm to the C6 proton at 6.05 ppm (1.3%) and a C12 proton at 3.45 ppm (3.3%). HPLC analysis by the previously described method showed one major peak with a purity of 99.1%.


- 54 -

Preparation of Example 6

3-((3S,6R,12aR)-6-Benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]pyrido[3,4-b]indole-3-yl)propionic acid




Example 6 was prepared from Example 5 by the following reaction.



Example 5

- 55 -



Example 6


To a mixture of Example 5 (1.13 g, 2.25 mmol) in acetic acid (11 mL) was added concentrated sulfuric acid (0.30 mL) dropwise. The resulting purple solution was stirred at room temperature for 30 minutes, after which water (50 mL) was added. The resulting white slurry was stirred at room temperature for 30 minutes. The solid was collected by vacuum filtration, and the solid residue was dissolved in a 70°C mixture of acetic acid and water (2:1, 50 mL). The yellow solution then was cooled to room temperature, then additional water (50 mL) was added. The solid was collected by vacuum filtration, and dried in a vacuum oven at 70°C overnight to yield Example 6 as a white solid (188 mg). The filtrate then was extracted with ethyl acetate to yield additional Example 6 (550 mg, total yield 74%): mp 221-231°C; ^1H NMR (300 MHz, DMSO- d_6): δ 11.05 (s, 1H), 8.62 (d, $J=3.8$ Hz, 1H), 7.52 (d, $J=7.4$ Hz, 1H), 7.29 (d, $J=7.4$ Hz, 1H), 7.12-6.92 (m, 2H), 6.88 (s, 1H), 6.78 (s, 2H), 6.09 (s, 1H), 5.92 (s, 2H), 4.44 (dd, $J=5.7, 4.4$ Hz, 1H), 3.92-3.79 (m,

- 56 -

1H), 3.58-3.42 (m, 1H), 3.02-2.88 (m, 1H), 2.35 (t, J=7.3 Hz, 2H), 2.05-1.82 (m, 2H); ¹³C NMR (125 MHz, DMSO-d₆): δ 173.7, 168.7, 167.7, 147.0, 146.0, 137.2, 136.1, 134.1, 125.7, 121.1, 119.0, 118.8, 118.0, 111.2, 108.0, 106.8, 104.5, 100.8, 55.5, 54.6, 54.3, 29.6, 28.4, 22.9 ppm; API MS m/z 448 [C₂₁H₂₀N₄O₃+H]⁺; [α]_D^{25°C}=+69.17° (c=0.5, DMSO). Anal. Calcd. for C₂₁H₂₀N₄O₃·0.75 H₂O: C, 62.64; H, 4.76; N, 9.13. Found: C, 62.55; H, 4.82; N, 9.00.


Preparation of Example 7

3-((3S,6R,12aR)-6-Benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]pyrido[3,4-b]indol-3-yl)-propionic acid isopropyl ester




Example 7 was prepared from Example 6 by the following reaction.

- 57 -



Example 6




Example 7

To a solution of Example 6 (558 mg, 1.25 mmol) and 4-(dimethylamino)pyridine (DMAP, 60 mg, 0.49 mmol) in isopropanol (1.0 mL, 13.10 mmol) and methylene chloride (10 mL), at 0°C under a nitrogen blanket, was added dicyclohexylcarbodiimide (DCC, 409 mg, 1.99 mmol) in one portion. The resulting white slurry was stirred at room temperature for 2 days. The white slurry was filtered under reduced pressure, then the solid was washed with methylene chloride (3 x 10 mL). The filtrate was concentrated



- 58 -

under reduced pressure and the residue was purified by flash column chromatography, eluting with methylene chloride/ethyl acetate (4:1), to provide Example 7 (450 mg), which contained residual dicyclohexylcarbodiimide. This solid was dissolved in methylene chloride (4 mL), and the solution was stored at -10°C overnight. The white crystals that formed were quickly filtered through a cold funnel under vacuum. The solid was washed with cold methylene chloride (2 x 1 mL), then dried in a vacuum oven at 60°C for 24 hours to provide Example 7 as a white solid (186 mg, 30%): TLC R_f (4:1 methylene chloride/ethyl acetate)=0.31; ¹H NMR (300 MHz, DMSO-d₆): δ 11.05 (s, 1H), 8.60 (d, J=3.7 Hz, 1H), 7.53 (d, J=7.4 Hz, 1H), 7.30 (d, J=7.4 Hz, 1H), 7.12-6.93 (m, 2H), 6.87 (s, 1H), 6.78 (s, 2H), 6.09 (s, 1H), 5.92 (s, 2H), 4.87 (sep, J=6.2 Hz, 1H), 4.44 (dd, J=5.7, 4.4 Hz, 1H), 3.90-3.79 (m, 1H), 3.55-3.47 (m, 1H), 3.02-2.87 (m, 1H), 2.45-2.30 (m, 2H), 2.03-1.87 (m, 2H), 1.22-1.05 (m, 6H); ¹³C NMR (125 MHz, DMSO-d₆): δ 171.6, 168.5, 167.7, 147.0, 146.0, 137.2, 136.1, 134.1, 125.7, 121.1, 119.0, 118.8, 118.0, 111.2, 108.0, 106.8, 104.5, 100.8, 67.1, 55.6, 54.6, 54.3, 29.8, 28.4, 22.9, 21.6 ppm; API MS m/z 491 [C₂₇H₂₇N₃O₆+H]⁺; [α]_D^{25°C}=+47.3° (c=0.5, CHCl₃). Anal. Calcd. for C₂₇H₂₇N₃O₆: C, 66.25; H, 5.56; N, 8.58. Found: C, 65.85; H, 5.63; N, 8.54. The stereochemistry was confirmed to be the *cis* isomer by an NOE difference experiment: positive NOE enhancements from the C12a proton at 4.44 ppm to the C6 proton at 6.09 ppm (1.8%) and a C12 proton at 3.50 ppm (2.7%).

- 59 -


Preparation of Example 8

The compound of Example 8 was prepared from Compound (III) and N-Cbz-(R)-(+)-oxazolidinecarboxylic acid as depicted in the following synthetic sequence.



Intermediate 10

- 60 -



Example 8

Preparation of Intermediate 10

To a solution of Cbz-(R)-(+) -oxazolidine-carboxylic acid (1.3 g, 5.1 mmol) in chloroform (20 mL), at 0°C, was added dicyclohexylcarbodiimide (1.1 g, 5.3 mmol) in one portion. The resulting white slurry was stirred at 0°C for 30 minutes, after which Compound (III) (1.7 g, 5.0 mmol) was added to the mixture. The resulting yellow slurry was warmed to room temperature, after which the slurry was heated at 60°C for 6 days. The solution then was cooled to room temperature, concentrated under reduced pressure to one-third of the slurry volume, and the resulting solid removed by vacuum filtration. The filtrate was concentrated under reduced pressure, and the residue was purified by flash column chromatography, eluting with methylene chloride/acetone (1:0 to 32:1), to provide Intermediate 10 as a colorless solid (0.56 g, 19%): TLC R_f (32:1 methylene chloride/acetone)=0.34.

- 61 -


Preparation of Example 8

(3R,6R,12aR)-6-Benzo[1,3]dioxo-5-yl-3-hydroxymethyl-2,3,6,7,12,12a-hexahydropyrazino[1',2':1,,6]pyrido[3,4-b]indole-1,4-dione


5

Intermediate 9 (0.55 g, 0.95 mmol) in methanol (50 mL) and acetic acid (15 mL) was treated with a catalytic amount of 10% palladium on carbon (0.08 g, about 10% wet). The mixture was stirred under a hydrogen atmosphere at 55°C for 3 hours, after which the palladium catalyst was removed by vacuum filtration through a plug of Celite, eluting with methanol. The filtrate was concentrated under reduced pressure to provide a yellow oil which was purified by flash column chromatography, eluting with methylene chloride/ethyl acetate/methanol (9:1:0.5), to yield Example 8 as a white foamy solid (0.17 g, 44%): mp 194-200°C; TLC R_f (8:1:0.5 methylene chloride/ethyl acetate/methanol)=0.43; ^1H NMR (300 MHz, CDCl_3): δ 7.87 (s, 1H), 7.58 (d, $J=6.6$ Hz, 1H), 7.29-7.25 (m, 2H), 7.20-7.12 (m, 2H), 6.82 (m, 1H), 6.73-6.64 (m, 3H), 6.16 (s, 1H), 5.86 (d, $J=5.7$ Hz, 2H), 4.39-4.33 (m, 1H), 4.15-4.12 (m, 1H), 4.02-3.90 (m, 2H), 3.75-3.69 (m, 1H), 3.28-3.19 (m, 1H); ^{13}C NMR (75 MHz, CDCl_3): δ 168.7, 148.6, 147.3, 136.7, 135.6, 132.9, 126.5, 122.8, 120.4, 120.1, 118.8, 111.4, 108.6, 107.1, 106.4, 101.3, 62.9, 56.9, 56.4, 55.6, 23.3 ppm; CI MS (methane) m/z 406 $[\text{C}_{22}\text{H}_{19}\text{N}_3\text{O}_5+\text{H}]^+$; $[\alpha]_{\text{D}}^{25^\circ\text{C}}=+58.7$ ($c=0.15$, methanol). Anal. Calcd. for $\text{C}_{23}\text{H}_{19}\text{N}_3\text{O}_5 \cdot 0.75 \text{H}_2\text{O}$: C, 63.08; H, 4.93; N, 10.03. Found: C, 63.17; H, 4.90; N, 9.79.


- 62 -

Preparation of Example 9

The compound of Example 9 can be prepared by the following synthetic sequence.




Intermediate 11




EDCl/Et₃N/CH₂Cl₂


- 63 -



Intermediate 12









Intermediate 13










Example 9









- 64 -

The following Examples 10-37 were prepared by methods analogous to Examples 1-9.








			
Example	X	R^2 1)	R^1
1		1	CH_3
2		1	H
3		1	H
4		1	CH_3
5		1	H

- 65 -








Example	X	R ² 1)	R ¹
6		1	H
7		1	H
8		1	H
9		1	CH ₃
10		1	CH ₃
11		1	H
12		1	H

Example	X	R ² 1)	R ¹
13		1	H
14		1	CH ₃
15		1	H
16		1	H
17		1	CH ₃
18		1	H
19		1	H
20		1	H




- 67 -

Example	X	R ² 1)	R ¹
21		1	H
22		1	H
23		1	H
24		1	H
25		1	H
26		1	H
27		1	H

- 68 -

Example	X	R ² 1)	R ¹
28		1	H
29		1	H
30		1	H
31		1	H
32		1	H
33		1	H
34		1	H


- 69 -

Example	X	R ² ¹⁾	R ¹
35		1	H
36		1	H
37		1	H

5

¹⁾ for R², the designation 1 is


10



15

and the designation 2 is

20



25

30

Compounds of the present invention can be formulated into tablets for oral administration. For example, a compound of formula (I) can be formed into a dispersion with a polymeric carrier by the

- 70 -

coprecipitation method set forth in WO 96/38131, incorporated herein by reference. The coprecipitated dispersion then can be blended with excipients, then pressed into tablets, which optionally are film-coated.

The compounds of structural formula (I) were tested for an ability to inhibit PDE5. The ability of a compound to inhibit PDE5 activity is related to the IC_{50} value for the compound, i.e., the concentration of inhibitor required for 50% inhibition of enzyme activity. The IC_{50} value for compounds of structural formula (I) were determined using recombinant human PDE5.

The compounds of the present invention typically exhibit an IC_{50} value against recombinant human PDE5 of less than about 50 μM , and preferably less than about 25 μM , and more preferably less than about 15 μM . The compounds of the present invention typically exhibit an IC_{50} value against recombinant human PDE5 of less than about 1 μM , and often less than about 0.05 μM . To achieve the full advantage of the present invention, a present PDE5 inhibitor has an IC_{50} of about 0.1 nM to about 15 μM .

The production of recombinant human PDEs and the IC_{50} determinations can be accomplished by well-known methods in the art. Exemplary methods are described as follows:

- 71 -

EXPRESSION OF HUMAN PDEs**Expression in *Saccharomyces cerevisiae* (Yeast)**

5 Recombinant production of human PDE1B,
PDE2, PDE4A, PDE4B, PDE4C, PDE4D, PDE5, and PDE7 was
carried out similarly to that described in Example 7
of U.S. Patent No. 5,702,936, incorporated herein by
reference, except that the yeast transformation
10 vector employed, which is derived from the basic
ADH2 plasmid described in Price et al., *Methods in*
Enzymology, 185, pp. 308-318 (1990), incorporated
yeast ADH2 promoter and terminator sequences and the
Saccharomyces cerevisiae host was the protease-defi-
15 cient strain BJ2-54 deposited on August 31, 1998
with the American Type Culture Collection, Manassas,
Virginia, under accession number ATCC 74465. Trans-
formed host cells were grown in 2X SC-leu medium, pH
6.2, with trace metals, and vitamins. After 24
20 hours, YEP medium-containing glycerol was added to a
final concentration of 2X YET/3% glycerol. Approxi-
mately 24 hr later, cells were harvested, washed,
and stored at -70°C.

HUMAN PHOSPHODIESTERASE PREPARATIONS**Phosphodiesterase Activity Determinations**

30 Phosphodiesterase activity of the prepara-
tions was determined as follows. PDE assays utiliz-
ing a charcoal separation technique were performed
essentially as described in Loughney et al. (1996).
In this assay, PDE activity converts [32P]cAMP or

- 72 -

[32P]cGMP to the corresponding [32P]5'-AMP or [32P]5'-GMP in proportion to the amount of PDE activity present. The [32P]5'-AMP or [32P]5'-GMP then was quantitatively converted to free [32P]phosphate and unlabeled adenosine or guanosine by the action of snake venom 5'-nucleotidase. Hence, the amount of [32P]phosphate liberated is proportional to enzyme activity. The assay was performed at 30°C in a 100 μ L reaction mixture containing (final concentrations) 40 mM Tris HCl (pH 8.0), 1 μ M ZnSO₄, 5 mM MgCl₂, and 0.1 mg/mL bovine serum albumin (BSA). PDE enzyme was present in quantities that yield <30% total hydrolysis of substrate (linear assay conditions). The assay was initiated by addition of substrate (1 mM [32P]cAMP or cGMP), and the mixture was incubated for 12 minutes. Seventy-five (75) μ g of *Crotalus atrox* venom then was added, and the incubation was continued for 3 minutes (15 minutes total). The reaction was stopped by addition of 200 μ L of activated charcoal (25 mg/mL suspension in 0.1 M NaH₂PO₄, pH 4). After centrifugation (750 X g for 3 minutes) to sediment the charcoal, a sample of the supernatant was taken for radioactivity determination in a scintillation counter and the PDE activity was calculated.

Purification of PDE5 from *S. cerevisiae*

Cell pellets (29 g) were thawed on ice with an equal volume of Lysis Buffer (25 mM Tris HCl, pH 8, 5 mM MgCl₂, 0.25 mM DTT, 1 mM benzamidine, and 10 μ M ZnSO₄). Cells were lysed in a Microfluidizer[®] (Microfluidics Corp.) using nitrogen at 20,000

- 73 -

psi. The lysate was centrifuged and filtered through 0.45 μ m disposable filters. The filtrate was applied to a 150 mL column of Q SEPHAROSE[®] Fast-Flow (Pharmacia). The column was washed with 1.5
5 volumes of Buffer A (20 mM Bis-Tris Propane, pH 6.8, 1 mM MgCl₂, 0.25 mM DTT, 10 μ M ZnSO₄) and eluted with a step gradient of 125 mM NaCl in Buffer A followed by a linear gradient of 125-1000 mM NaCl in Buffer A. Active fractions from the linear gradient were
10 applied to a 180 mL hydroxyapatite column in Buffer B (20 mM Bis-Tris Propane (pH 6.8), 1 mM MgCl₂, 0.25 mM DTT, 10 μ M ZnSO₄, and 250 mM KCl). After loading, the column was washed with 2 volumes of Buffer B and eluted with a linear gradient of 0-125 mM
15 potassium phosphate in Buffer B. Active fractions were pooled, precipitated with 60% ammonium sulfate, and resuspended in Buffer C (20 mM Bis-Tris Propane, pH 6.8, 125 mM NaCl, 0.5 mM DTT, and 10 μ M ZnSO₄). The pool was applied to a 140 mL column of
20 SEPHACRYL[®] S-300 HR and eluted with Buffer C. Active fractions were diluted to 50% glycerol and stored at -20°C.

The resultant preparations were about 85% pure by SDS-PAGE. These preparations had specific
25 activities of about 3 μ mol cGMP hydrolyzed per minute per milligram protein.

- 74 -

Inhibitory Effect on cGMP-PDE

cGMP-PDE activity of compounds of the present invention was measured using a one-step assay adapted from Wells et al., *Biochim. Biophys. Acta*, 384, 430 (1975). The reaction medium contained 50 mM Tris-HCl, pH 7.5, 5 mM magnesium acetate, 250 μ g/ml 5'-Nucleotidase, 1 mM EGTA, and 0.15 μ M 8-[H³]-cGMP. Unless otherwise indicated, the enzyme used was a human recombinant PDE5 (ICOS Corp., Bothell, Washington).

Compounds of the invention were dissolved in DMSO finally present at 2% in the assay. The incubation time was 30 minutes during which the total substrate conversion did not exceed 30%.

The IC₅₀ values for the compounds examined were determined from concentration-response curves typically using concentrations ranging from 10 nM to 10 μ M. Tests against other PDE enzymes using standard methodology showed that compounds of the invention are selective for the cGMP-specific PDE enzyme.

Biological Data

The compounds according to the present invention were typically found to exhibit an IC₅₀ value of less than 500 nM (i.e., 0.5 μ M). An *in vitro* test data for representative compounds of the invention is given in the following table

- 75 -

Table 1: <i>In vitro</i> Results	
Example	PDE5 IC ₅₀ (nM)
1	161.0
2	1.2
3	113.0
4	593.0
5	0.6
6	1.0
7	0.2
8	23.0
12	0.03
15	8.6
16	9.5
18	9.2
19	20
20	600
21	420
22	80
23	10
24	90
25	80
26	50
27	90
28	10
29	30
30	8.6
31	7.8
32	7.0
33	9.0
34	7.5
35	20
36	7.7

- 76 -

Table 1: <i>In vitro</i> Results	
Example	PDE5 IC ₅₀ (nM)
37	8.0

5


Obviously, many modifications and variations of the invention as hereinbefore set forth can be made without departing from the spirit and scope thereof, and, therefore, only such limitations should be imposed as are indicated by the appended claims.

10

- 77 -

WHAT IS CLAIMED IS:


1. A compound having a formula



wherein R^0 , independently, is selected from the group consisting of halo and C_{1-6} alkyl;

R^1 is selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo C_{1-6} -alkyl, C_{3-8} cycloalkyl, C_{3-8} cycloalkyl C_{1-3} alkyl, aryl C_{1-3} -alkyl, and heteroaryl C_{1-3} alkyl;

R^2 is selected from the group consisting of an optionally substituted monocyclic aromatic ring selected from the group consisting of benzene, thiophene, furan, and pyridine, and an optionally substituted bicyclic ring



wherein the fused ring A is a 5- or 6-membered ring, saturated or partially or fully unsaturated, and comprises carbon atoms and optionally one or two heteroatoms selected from oxygen, sulfur, and nitrogen;

- 78 -

X is selected from the group consisting of NR^a , O, S, CR^3R^4 , $\text{CH}_2\text{CR}^3\text{R}^5$, and $\text{CR}^3\text{R}^5\text{CH}_2$;

R^3 is selected from the group consisting of hydrogen and C_{1-3} alkyl;

or R^1 and R^3 together represent a 3- or 4-membered alkyl or alkenyl chain component of a 5- or 6-membered ring,

or NR^a , R^1 , and the nitrogen to which R^1 is attached, form a 5- or 6-membered heterocyclic ring;

R^4 is selected from the group consisting of C_{1-6} alkyl, haloC_{1-4} alkyl, aryl, heteroaryl, Het, C_{3-8} -cycloalkyl, OR^a , $\text{C}(=\text{O})\text{OR}^a$, $\text{C}(=\text{O})\text{R}^a$, $\text{C}(=\text{O})\text{NR}^a\text{SO}_2\text{R}^b$, $\text{C}(=\text{O})\text{NR}^a\text{R}^b$, $\text{C}(=\text{S})\text{NR}^a\text{R}^b$, C_{1-6} alkylene $\text{C}(=\text{O})\text{OR}^a$, C_{1-4} alkylene $\text{C}(=\text{O})\text{NR}^a\text{R}^b$, C_{1-4} alkylene NR^aR^b , C_{1-4} alkylene OR^a , C_{1-4} alkylene $\text{SO}_2\text{NR}^a\text{R}^b$, C_{1-4} alkylenearyl, C_{1-4} alkyleneheteroaryl, C_{1-4} alkyleneHet, C_{1-4} alkylene $\text{C}(=\text{O})\text{C}_{1-4}$ alkyleneHet, C_{1-4} alkylene $\text{C}(=\text{O})\text{Het}$, C_{1-4} alkylene $\text{C}(=\text{O})\text{OR}^a$, $\text{C}(=\text{O})\text{C}_{1-4}$ alkyleneHet, $\text{C}(=\text{O})\text{NR}^a\text{alkyleneOR}^b$, $\text{C}(=\text{O})\text{NR}^a\text{C}_{1-4}$ alkyleneHet, $\text{NR}^a\text{C}_{1-4}$ alkylene NR^aR^b , $\text{NR}^a\text{C}(=\text{O})\text{R}^b$, $\text{NR}^a\text{C}(=\text{O})\text{NR}^a\text{R}^b$, $\text{N}(\text{SO}_2\text{C}_{1-4}\text{alkyl})_2$, and $\text{NR}^a(\text{SO}_2\text{C}_{1-4}\text{alkyl})$, with the proviso that when R^3 is hydrogen, R^4 is different from C_{1-3} alkyl;

or R^1 and R^4 together represent a 3- or 4-membered carbocyclic or heterocyclic chain component, either saturated or unsaturated, of a 5- or 6-membered ring;

or R^3 and R^4 together represent a 4- to 6-membered alkyl or alkenyl chain component of a 5- to 7-membered ring;

R^5 is selected from the group consisting of hydrogen, C_{1-6} alkyl, haloC_{1-4} alkyl, aryl, heteroaryl, Het, C_{3-8} cycloalkyl, OR^a , $\text{C}(=\text{O})\text{OR}^a$, $\text{C}(=\text{O})\text{R}^a$, $\text{C}(=\text{O})\text{NR}^a\text{SO}_2\text{R}^b$, $\text{C}(=\text{O})\text{NR}^a\text{R}^b$, $\text{C}(=\text{S})\text{NR}^a\text{R}^b$, C_{1-6} alkylene $\text{C}(=\text{O})\text{OR}^a$,

- 79 -

C_{1-4} alkyleneC(=O)NR^aR^b, C_{1-4} alkyleneNR^aR^b, C_{1-4} alkylene-OR^a, C_{1-4} alkyleneSO₂NR^aR^b, C_{1-4} alkylenearyl, C_{1-4} alkyleneheteroaryl, C_{1-4} alkyleneHet, C_{1-4} alkyleneC(=O) C_{1-4} alkyleneHet, C_{1-4} alkyleneC(=O)Het, C_{1-4} alkyleneC(=O)-OR^a, C(=O) C_{1-4} alkyleneHet, C(=O)NR^aalkyleneOR^b, C(=O)-NR^a C_{1-4} alkyleneHet, NR^a C_{1-4} alkyleneNR^aR^b, NR^aC(=O)R^b, NR^aC(=O)NR^aR^b, N(SO₂ C_{1-4} alkyl)₂, and NR^a(SO₂ C_{1-4} alkyl);

R^a and R^b, independently, are selected from the group consisting of hydrogen, C_{1-8} alkyl, aryl, heteroaryl, halo C_{1-6} alkyl, C_{1-4} alkyleneHet, aryl C_{1-3} alkyl, heteroaryl C_{1-3} alkyl, C_{1-4} alkylenearyl, C_{3-8} cycloalkyl, C_{1-3} alkyleneheteroaryl, and Het;


Het is a 4- to 7-membered heterocyclic group, saturated or partially unsaturated, containing at least one heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur, and optionally substituted with one or more C_{1-6} alkyl, NR^aR^b, and C(=O)OR^a;

q is 0, 1, 2, 3, or 4; and

pharmaceutically acceptable salts and solvates thereof.

- 80 -

2. The compound of claim 1 represented by the formula



and pharmaceutically acceptable salts and solvates thereof.

3. The compound of claim 1 wherein R^4 and R^5 are selected from the group consisting of C_{1-6} alkyl, C_{1-6} alkyleneC(=O)OR^a, C(=O)OR^a, C(=S)NR^aR^b, C_{5-7} spiro when combined with R^3 , C_{1-4} alkyleneC(=O)NR^aR^b, C_{1-4} alkyleneNR^aR^b, C_{1-4} alkyleneOR^a, C_{1-4} alkyleneSO₂NR^aR^b, C_{1-4} alkyleneheteroaryl, haloC₁₋₄alkyl, C_{1-4} alkyleneHet, C_{1-4} alkyleneC(=O) C_{1-4} alkyleneHet, and C_{1-4} alkylenearyl; and R^a and R^b , independently, are selected from the group consisting of C_{1-8} alkyl, hydrogen, C_{1-4} alkylenearyl, Het, C_{1-4} alkyleneHet, C_{3-8} cycloalkyl, and haloC₁₋₆alkyl.





4. The compound of claim 3 wherein Het, aryl, and heteroaryl are optionally substituted with one or more of C(=O)OR^a, NR^aR^b, and C_{1-6} alkyl.

5. The compound of claim 1 wherein X is selected from the group consisting of CH₂CH₂, NH, and NCH₃.



- 81 -

6. The compound of claim 1 wherein R^3 is selected from the group consisting of hydrogen and methyl.


7. The compound of claim 1 wherein R^4 is selected from the group consisting of C_{1-4} alkyl, C_{1-4} alkyleneC(=O)NH₂, C_{1-4} alkyleneNH₂, C_{1-4} alkyleneC(=O)OC₁₋₇alkyl, C_{1-6} alkyleneC(=O)OR^a, C_{1-3} alkyleneOH, C_{1-4} alkyleneSO₂NH₂, C_{1-4} alkyleneOC₁₋₃alkylenearyl, halo- C_{1-3} alkyl, C_{1-3} alkylenearyl, C_{1-4} alkyleneC(=O)OC₁₋₃halo-alkyl,



- 82 -




, and



8. The compound of claim 1 wherein R^3 and R^4 are taken together to form a 5- or 6-membered spiro group.


9. The compound of claim 1 wherein R^2 is




wherein n is an integer 1 or 2, and Y and Z , independently, are CH_2 , O , S , or NR^a .

10. The compound of claim 1 wherein R^2 , substituted or unsubstituted, is selected from the group consisting of


- 83 -



,






, and






.

- 84 -




11. A compound of claim 1 selected from the group consisting of







- 85 -





- 86 -









and pharmaceutically acceptable salts and solvates thereof.

- 90 -

12. A compound of claim 1 selected from the group consisting of

(3S,6R,12aR)-3-(4-aminobutyl)-6-benzo[1,3]dioxol-5-yl-2,3,6,7,12,12a-hexahydropyrazino[1',2':1,6]-pyrido[3,4-b]indole-1,4-dione,

((3S,6R,12aR)-6-benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]-pyrido[3,4-b]indol-3-yl)acetic acid tert-butyl ester,

(3S,6R,12aR)-6-benzo[1,3]dioxol-5-yl-3-benzyloxy-methyl-2,3,6,7,12,12a-hexahydropyrazino[1',2':1,6]-pyrido[3,4-b]indole-1,4-dione,

((3S,6R,12aR)-6-benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]-pyrido[3,4-b]indol-3-yl)acetic acid,

((3R,6R,12aR)-6-benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]-pyrido[3,4-b]indol-3-yl)acetic acid,

(6R,12aR)-6-benzo[1,3]dioxol-5-yl-3-pyrazol-1-ylmethyl-2,3,6,7,12,12a-hexahydropyrazino[1',2':1,6]pyrido[3,4-b]indole-1,4-dione,

(3S,6R,12aR)-3-(2-aminoethyl)-6-benzo[1,3]dioxol-5-yl-2,3,6,7,12,12a-hexahydropyrazino[1',2':1,6]-pyrido[3,4-b]indole-1,4-dione,

(3S,6R,12aR)-3-aminomethyl-6-benzo[1,3]dioxol-5-yl-2,3,6,7,12,12a-hexahydropyrazino[1',2':1,6]-pyrido[3,4-b]indole-1,4-dione,

(3R,6R,12aR)-6-benzo[1,3]dioxol-5-yl-3-chloromethyl-2,3,6,7,12,12a-hexahydropyrazino[1',2':1,6]-pyrido[3,4-b]indole-1,4-dione

2-((3S,6R,12aR)-6-benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]-

- 91 -

pyrido[3,4-b]indol-3-yl)-N-(4-dimethylaminobenzyl)-acetamide,
(3S,6R,12aR)-6-benzo[1,3]dioxol-5-yl-3-[2-(4-methylpiperazin-1-yl)-2-oxo-ethyl]-2,3,6,7,12,12a-hexahydropyrazino[1',2':1,6]pyrido[3,4-b]indole-1,4-dione,
2-((6R,12aR)-6-benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]pyrido[3,4-b]indol-3-yl)-N-(2-pyrrolidin-1-ylethyl)acetamide,
(3S,6R,12aR)-6-benzo[1,3]dioxol-5-yl-3-pyridin-3-ylmethyl-2,3,6,7,12,12a-hexahydropyrazino[1',2':1,6]pyrido[3,4-b]indole-1,4-dione,
((3S,6R,12aR)-6-benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]pyrido[3,4-b]indol-3-yl)acetic acid methyl ester,
((3S,6R,12aR)-6-benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]pyrido[3,4-b]indol-3-yl)acetic acid heptyl ester,
((3S,6R,12aR)-6-benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]pyrido[3,4-b]indol-3-yl)acetic acid ethyl ester,
((3S,6R,12aR)-6-benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]pyrido[3,4-b]indol-3-yl)acetic acid isopropyl ester,
((3S,6R,12aR)-6-benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]pyrido[3,4-b]indol-3-yl)acetic acid cyclopentyl ester,
((3S,6R,12aR)-6-benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]pyrido[3,4-b]indol-3-yl)acetic acid 2,2,2-trifluoroethyl ester,

- 92 -

3-((3S,6R,12aR)-6-benzofuran-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]-pyrido[3,4-b]indol-3-yl)propionic acid tert-butyl ester,


((3S,6R,12aR)-6-benzofuran-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]-pyrido[3,4-b]indol-3-yl)acetic acid tert-butyl ester,

3-((3S,6R,12aR)-6-benzo[1,3]dioxol-5-yl-1,4-dioxo-1,2,3,4,6,7,12,12a-octahydropyrazino[1',2':1,6]-pyrido[3,4-b]indol-3-yl)propionic acid ethyl ester,


and pharmaceutically acceptable salts and solvates thereof.

- 93 -


13. A compound having a formula



wherein R¹ is hydrogen or methyl, R² is







or















, and













X is selected from the group consisting of













- 94 -



- 95 -



- 96 -



14. A pharmaceutical composition comprising a compound of claim 1, together with a pharmaceutically acceptable diluent or carrier.

- 97 -

15. A method of treating a male or female animal in the treatment of a condition where inhibition of a cGMP-specific PDE is of a therapeutic benefit comprising treating said animal with an effective amount of a pharmaceutical composition comprising a compound of claim 1, together with a pharmaceutically acceptable diluent or carrier.

16. The method of claim 15 wherein the condition is male erectile dysfunction.

17. The method of claim 16 wherein the treatment is an oral treatment.

18. The method of claim 15 wherein the condition is female arousal disorder.

19. The method of claim 18 wherein the treatment is an oral treatment.

- 98 -

20. The method of claim 15 wherein the condition is selected from the group consisting of stable angina, unstable angina, variant angina, hypertension, pulmonary hypertension, chronic obstructive pulmonary disease, malignant hypertension, pheochromocytoma, acute respiratory distress syndrome, congestive heart failure, acute renal failure, chronic renal failure, atherosclerosis, a condition of reduced blood vessel patency, a peripheral vascular disease, a vascular disorder, thrombocythemia, an inflammatory disease, myocardial infarction, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma, peptic ulcer, a gut motility disorder, postpercutaneous transluminal coronary angioplasty, carotid angioplasty, post-bypass surgery graft stenosis, osteoporosis, preterm labor, benign prostatic hypertrophy, and irritable bowel syndrome.

21. A method of treating a condition where inhibition of a cGMP-specific PDE is of therapeutic benefit, in a human or a nonhuman animal body, comprising administering to said body a therapeutically effective amount of a compound of claim 1.

22. A method for the curative or prophylactic treatment of male erectile dysfunction or female arousal disorder, comprising administration of an effective dose of a compound of claim 1, and pharmaceutically acceptable salts and solvates thereof, to an animal.

- 99 -

23. Use of a compound of claim 1 for the manufacture of a medicament for the curative or prophylactic treatment of a condition where inhibition of a cGMP-specific PDE is of a therapeutic benefit.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 01/21678

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D471/16 A61K31/495 C07D471/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 03985 A (GLAXO WELLCOME LAB SA ;DAUGAN ALAIN CLAUDE MARIE (FR); GELLIBERT F) 6 February 1997 (1997-02-06) page 1, line 1-9; claim 1 ----	1-23
X	US 6 046 199 A (PAMUKCU RIFAT ET AL) 4 April 2000 (2000-04-04) claim 1 -----	1-23
X	WO 97 03675 A (GLAXO WELLCOME LAB SA ;DAUGAN ALAIN CLAUDE MARIE (FR)) 6 February 1997 (1997-02-06) page 1, line 1-12; claim 1 -----	1-23
X	WO 95 19978 A (GLAXO LAB SA ;DAUGAN ALAIN CLAUDE MARIE (FR)) 27 July 1995 (1995-07-27) page 1, line 1-9; claim 1 ----- -/-	1-23

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

20 December 2001

Date of mailing of the international search report

16/01/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Samsam Bakhtiary, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/21678

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 1 022 026 A (PFIZER LTD ; PFIZER RES & DEV (IE)) 26 July 2000 (2000-07-26) page 8, line 22-57 ----	1-23
X	SYBERTZ E ET AL: "Inhibitors of PDE1 and PDE5 cGMP phosphodiesterases: patents and therapeutic potential" EXPERT OPINION ON THERAPEUTIC PATENTS, ASHLEY PUBLICATIONS, GB, vol. 7, no. 6, 1997, pages 631-639, XP002178692 ISSN: 1354-3776 page 635; example 15 ----	1-23
P,X	WO 00 66114 A (PULLMAN WILLIAM E ; LILLY ICOS LLC (US); ALLEMEIER LORA L (US); BRA) 9 November 2000 (2000-11-09) page 11, line 1-10 ----	1-23
P,X	WO 00 66099 A (LILLY ICOS LLC ; PULLMAN WILLIAM ERNEST (US); WHITAKER JOHN STEVEN) 9 November 2000 (2000-11-09) page 1, line 10-25; claim 1 -----	1-23

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claims 15-22 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.1

Claims Nos.: 15-22

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

INTERNATIONAL SEARCH REPORT

Int. Patent Application No.

PCT/US 01/21678

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 9703985	A	06-02-1997	AU	702324 B2	18-02-1999
			AU	6419296 A	18-02-1997
			BR	9609780 A	09-03-1999
			CA	2226761 A1	06-02-1997
			CN	1195349 A ,B	07-10-1998
			CZ	9800032 A3	17-06-1998
			WO	9703985 A1	06-02-1997
			EP	0846118 A1	10-06-1998
			HR	960321 A1	31-08-1998
			HU	9900006 A2	28-04-1999
			JP	11509535 T	24-08-1999
			NO	980154 A	10-03-1998
			PL	324527 A1	08-06-1998
			SK	3898 A3	04-11-1998
			US	5981527 A	09-11-1999
			US	6143746 A	07-11-2000
<hr/>					
US 6046199	A	04-04-2000	NONE		
<hr/>					
WO 9703675	A	06-02-1997	AU	704955 B2	13-05-1999
			AU	6419196 A	18-02-1997
			BR	9609758 A	26-01-1999
			CA	2226784 A1	06-02-1997
			CN	1195290 A	07-10-1998
			CZ	9800033 A3	13-05-1998
			WO	9703675 A1	06-02-1997
			EP	0839040 A1	06-05-1998
			HU	9900065 A2	28-05-1999
			JP	11509221 T	17-08-1999
			NO	980153 A	10-03-1998
			PL	324495 A1	25-05-1998
			SK	3998 A3	08-07-1998
			US	6140329 A	31-10-2000
<hr/>					
WO 9519978	A	27-07-1995	AP	556 A	07-11-1996
			AT	169018 T	15-08-1998
			AU	689205 B2	26-03-1998
			AU	1574895 A	08-08-1995
			AU	707055 B2	01-07-1999
			AU	7391298 A	20-08-1998
			BG	62733 B1	30-06-2000
			BG	100727 A	28-02-1997
			BR	9506559 A	28-10-1997
			CA	2181377 A1	27-07-1995
			CN	1143963 A ,B	26-02-1997
			CN	1224720 A ,B	04-08-1999
			CZ	9602116 A3	11-06-1997
			DE	69503753 D1	03-09-1998
			DE	69503753 T2	21-01-1999
			DK	740668 T3	03-05-1999
			EE	3231 B1	16-12-1996
			WO	9519978 A1	27-07-1995
			EP	0740668 A1	06-11-1996
			ES	2122543 T3	16-12-1998
			FI	962927 A	19-07-1996
			HK	1013286 A1	19-05-2000
			HR	950023 A1	30-04-1998
			HU	74943 A2	28-03-1997

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/21678

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9519978	A	IL 112384 A	16-08-1998
		JP 9508113 T	19-08-1997
		LV 11690 A	20-02-1997
		LV 11690 B	20-06-1997
		NO 963015 A	09-09-1996
		NZ 279199 A	26-01-1998
		PL 315559 A1	12-11-1996
		RU 2142463 C1	10-12-1999
		SG 49184 A1	18-05-1998
		SI 740668 T1	28-02-1999
		SK 94096 A3	09-04-1997
		TW 378210 B	01-01-2000
		US 6025494 A	15-02-2000
		US 6143746 A	07-11-2000
		US 6127542 A	03-10-2000
		US 5859006 A	12-01-1999
		ZA 9500424 A	27-09-1995
EP 1022026	A	26-07-2000	
		AU 6178899 A	01-06-2000
		EP 1022026 A2	26-07-2000
		HU 9904434 A2	28-08-2000
		JP 2000159672 A	13-06-2000
		US 6225315 B1	01-05-2001
WO 0066114	A	09-11-2000	
		AU 4663100 A	17-11-2000
		WO 0066114 A1	09-11-2000
WO 0066099	A	09-11-2000	
		AU 3012900 A	02-11-2000
		AU 4490800 A	17-11-2000
		CN 1292264 A	25-04-2001
		DE 10021266 A1	16-11-2000
		DE 20007861 U1	24-08-2000
		DK 200000677 A	31-10-2000
		FI 20000976 A	30-10-2000
		FR 2795646 A1	05-01-2001
		GB 2351663 A	10-01-2001
		GR 2000100153 A	29-12-2000
		HU 0001632 A2	28-05-2001
		JP 2000336043 A	05-12-2000
		LT 2000035 A ,B	27-11-2000
		LV 12560 A	20-11-2000
		LV 12560 B	20-04-2001
		NL 1015027 C2	14-02-2001
		NL 1015027 A1	31-10-2000
		NO 20002097 A	26-10-2001
		PL 339897 A1	06-11-2000
		PT 102457 A	30-11-2000
		SE 0001518 A	31-10-2000
		SI 20361 A	30-04-2001
		TR 200001132 A2	22-01-2001
		WO 0066099 A2	09-11-2000